

Lebanon



BROADLEAF WEED KILLER

with **TRIMEC[®] HERBICIDE**



CONTROLS: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds, as listed on the back panel.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid.....	1.37%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid.....	0.31%
Dicamba: 3,6-dichloro-o-anisic acid.....	0.13%
OTHER INGREDIENTS:	98.19%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 27.4 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton.
 - 6.2 lb. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
 - 2.6 lb. 3,6-dichloro-o-anisic acid equivalent per ton.
- Isomer Specific by AOAC Methods

EPA REG. NO. 2217-640-961

(L) (D)
EPA Est. 961-PA-01; 961-IL-01
Superscript used is first letter of lot number.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.




LebanonTurf

Distributed by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
1-800-233-0628 • (717) 273-1685
Manufactured in U.S.A.

24-35799

NET WT. 20 LBS. (9.07 KG)
Covers up to 10,000 sq. ft.



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

FIRST AID	
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- For residential use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Apply when the air is calm. Do not apply to newly seeded lawns. Do not exceed specified dosages for any area. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Lebanon Broadleaf Weed Killer with Trimec® Herbicide controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others as listed which are commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

For use on established turf of warm season grasses including bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses including Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq. ft.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

SUGGESTED SPREADER SETTINGS			
Spreader	2#/1,000 sq. ft.	Spreader	2#/1,000 sq. ft.
LebanonTurf	4	Lesco (letter/Numeric)	G/15
Andersons AccuPro	H	PennMulch HVO	J
Earthway Rotary	16	ProScape SS	J
Gandy	24	Spyker	4
Lely	4II	Vicon (all models)	22
The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 pound of product per 500 square feet.			

Weeds

Annual yellow sweetclover	Buckhorn	Clover, white	Hawkweed	Mustard	Prostrate vervain	Spotted catsear	Wild four-o'clock
Aster	Bulbous buttercup	Cockle	Healall	Narrowleaf plantain	Puncturevine	Spotted spurge	Wild garlic
Austrian fieldcress	Bull thistle	Cocklebur	Heartleaf drymary	Narrowleaf vetch	Purslane, common	Spurweed	Wild geranium
Bedstraw	Bullnettle	Common mullein	Heath aster	Nettle	Ragweed	Stinging nettle	Wild lettuce
Beggarticks	Burclover	Creeping jenny	Hedge bindweed	Orange hawkweed	Red sorrel	Strawberry, India mock	Wild marigold
Betony, Florida	Burdock	Cudweed	Hedge mustard	Oriental cocklebur	Redroot pigweed	Tall nettle	Wild mustard
Bindweed, field	Burning nettle	Curly dock	Hemp	Oxalis	Redstem filaree	Tall vervain	Wild onion
Bird vetch	Burweed	Daisy, English	Henbit	Parsley-piert	Rough cinquefoil	Tansy ragwort	Wild parsnip
Bitter wintercress	Buttercup	Daisy fleabane	Hoary cress	Parsnip	Rough fleabane	Tansy mustard	Wild radish
Bittercress, hairy	Buttonweed	Daisy, oxeye	Hoary plantain	Pearlwort	Russian pigweed	Tanweed	Wild rape
Bitterweed	Canada thistle	Dandelion	Hoary vervain	Pennycress	Russian thistle	Thistle	Wild strawberry
Black-eyed Susan	Carolina geranium	Dichondra	Horsenettle	Pepperweed	Scarlet pimpernel	Trailing crownvetch	Wild sweet potato
Black medic	Carpetweed	Dogbane	Jimsonweed	Pigweed	Scotch thistle	Tumble mustard	Wild vetch
Black mustard	Catchweed bedstraw	Dogfennel	Knawel	Pineywoods bedstraw	Sheep sorrel	Tumble pigweed	Woodsorrel
Blackseed plantain	Catnip	Dollarweed	Knotweed	Plains coreopsis	Shepherdspurge	Velvetleaf	Woolly croton
Blessed thistle	Catsear	False dandelion	Kochia	Plantain	Slender plantain	Venice mallow	Woolly morningglory
Bloodflower milkweed	Chickweed, common	Falseflax	Lambsquarters	Poison ivy	Smallflower galinsoga	Virginia buttonweed	Woolly plantain
Blue lettuce	Chickweed, mouseear	False sunflower	Lespedeza	Poison oak	Smooth dock	Virginia creeper	Wormseed
Blue vervain	Chicory	Fiddleneck	Mallow	Pokeweed	Smooth pigweed	Virginia pepperweed	Yarrow
Bracted plantain	Cinquefoil	Florida pusley	Matchweed	Poorjoe	Sorrel	Wavyleaf bullthistle	Yellow rocket
Brassbuttons	Clover, crimson	Frenchweed	Mexicanweed	Prairie sunflower	Sowthistle	Western clematis	Yellowflower
Bristly oxtongue	Clover, hop	Galinsoga	Milk vetch	Prickly lettuce	Spanishneedles	Western salsify	pepperweed
Broadleaf dock	Clover, red	Goldenrod	Morningglory	Prickly sida	Speedwell	White mustard	
Broadleaf plantain	Clover, strawberry	Ground ivy	Mouseear hawkweed	Prostrate knotweed	Spiny amaranth	Wild aster	
Broomweed	Clover, sweet	Gumweed	Mugwort	Prostrate pigweed	Spiny cocklebur	Wild buckwheat	
		Hairy fleabane	Musk thistle	Prostrate spurge	Spiny sowthistle	Wild carrot	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.