

GREENVIEW®

BROADLEAF WEED CONTROL PLUS LAWN FOOD



Helps lawns retain water to
protect against heat & drought



Speeds root growth to
improve nutrient absorption

Feeds your lawn up to 8 weeks

Apply when listed weeds are
growing in spring or early fall



For more information, visit www.greenviewfertilizer.com

**Kills Over 250
Weeds as listed**



**NET WEIGHT 39 lbs. (17.69 kg)
Covers 15,000 sq. ft.**

ACTIVE INGREDIENTS:
2,4-D, 2-ethylhexyl ester..... 1.108%
Mecoprop-p acid..... 0.167%
Dicamba acid..... 0.071%
OTHER INGREDIENTS:..... 98.654%
TOTAL :.....100.000%

THIS PRODUCT CONTAINS:
14.70 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
3.34 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
1.42 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9

Isomer Specific by AOAC Methods
TRIMEC is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panel for First Aid and
additional Precautionary Statements.

**GREENVIEW® BROADLEAF WEED CONTROL
PLUS LAWN FOOD**



Poor soil may have fewer of the nutrients needed to promote a vigorous lawn. GreenSmart[†] works by enriching your soil with nutrients.

GreenSmart[†] revitalizes the soil, which aids in the development of a larger, deeper and better root system.

The enhanced habitat is proven to produce a green, thick, vigorous lawn and reduces the need for watering.

[†] Enhanced Efficiency Fertilizer

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
IF IN EYES:	•Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Treatment Information call (888)208-1368.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
•Do not apply this product in a way that will contact any person or pet either directly or through drift.
•This product is for residential use.
•Do not allow people (other than applicator) or pets on treatment area during treatment. Do not enter treatment area until dust has settled.

Application Instructions

GreenView Broadleaf Weed Control plus Lawn Food contains Trimec[®] Herbicide. Trimec[®] controls a wide range of lawn weeds, including dandelion, chickweed, plantain, henbit, spurge and many other listed weeds commonly found in home lawns. In addition, a 27-0-4 fertilizer provides a supply of lawn food. Possible retreatment may be necessary if (a) the lawn is heavily overrun by weeds or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made after 30 days. Treated areas may be seeded 3 weeks after application.

USE PRECAUTIONS AND RESTRICTIONS:

Do not use on dichondra, carpetgrass, 'Floritam' St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.

Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of or over surface roots of trees, shrubs, and other ornamentals.

Reseeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY:

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made after 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

Maximum Use Rate Statement

Do not apply more than 2.6 pounds of product per 1000 square feet of turfgrass per application. If necessary, a second application may be made no sooner than 30 days after the first. Do not apply more than 5.2 pounds of product per 1000 square feet per year.

The maximum broadcast application rate per acre is 1.28 lbs. 2,4-D ae, 0.29 lb. MCPP-p ae and 0.12 lb. dicamba ae per acre per application. The maximum seasonal rate is 2.56 lbs. 2,4-D ae, 0.58 lb. MCPP-p ae and 0.25 lb. dicamba ae per acre.

WHEN TO APPLY:

Apply when listed broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

APPLICATION INSTRUCTIONS:


1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when the dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

GreenView Broadleaf Weed Control plus Lawn Food Kills more than 250 weeds as listed!

Amaranth, purple	Cocklebur, spiny	Matchweed	Ragweed, Western	Wild garlic
Amaranth, slender	Common mullein	Mexicanweed	Redstem filaree	Wild lettuce
Aster, heath	Corn Spurry	Moneywort	Rough cinquefoil	Wild marigold
Aster, white heath	Creeping Beggarweed	Morningglory, bigroot	Scarlet pimpernel	Wild mustard
Aster, purple	Creeping Jenny	Morningglory, ivyleaf	Shepard's purse	(yellow rocket)
Aster, white prairie	(creeping Charlie)	Morningglory, Mexican	Smartweed, ladysthumb	Wild onion
Artichoke	Cupid shaving brush	Morningglory, red	Smartweed, pale	Wild pansnip
Austrian fieldcress	Cudweed	Morningglory, tall	Smartweed, Pennsylvania	Wild radish
Bedstraw, Catchweed	Daisy, English	Morningglory, woolly	Smooth chaff-flower	Wild rape
Bedstraw, smooth	Daisy, oxeye	Mugwort	Smoothdock	Wild strawberry
Bedstraw, Northern	Dandelion, common	Mustard, ball	Sorrel, red (sheep)	Wild sweet potato
Bedstraw, pinewoods	Dandelion, false	Mustard, black	Sowthistle, annual	Wild violet
Beggarticks, hairy	Day flower	Mustard, blue	Sowthistle, spiny	Woodsorrel, creeping
Betony, Florida	Deadnettle	Mustard, hedge	Spanish needles	Woodsorrel, violet
Bindweed, Field	Dichondra	Mustard, Indian	Speedwell, bi-lobed	Woodsorrel, yellow
Bindweed, Hedge	Dock, broadleaf	Mustard, tansy	Speedwell, common	(Oxalis)
Bitter Wintercress	Dock, curly	Mustard, tumble	Speedwell, corn	Woolly croton
Bittercress, hairy	Dogbane	Mustard, white	Speedwell, ivyleaf	Wormseed
Bitterweed	Dogfennel	Mustard, wormseed	Speedwell, Persian	Yarrow, common
Black-eyed Susan	Dollarweed (Pennywort)	Nettle, stinging (burning)	Speedwell, purslane	Yarrow, Western
Black Medic	Dove weed	Nettle, tall	Speedwell, slender	Yarrow, yellow
Blood flower milkweed	Elderberry	Nettle, wood	Speedwell, snow	
Blue lettuce	Falseflax	Parsley-piert	Speedwell, thymeleaf	
Brassbuttons	False sunflower	Parsnip	Spiny amaranth	
Bristly oxtongue	Fiddleneck	Pearlwort	Spurge, spotted	
Broomweed	Filaree, redstem	Pennycress (frenchweed)	Spurge, prostrate	
Buckhorn	Filaree, whitestem	Pepperweed, perennial	Spurweed	
Bullnettle	Fleabane, daisy (annual)	Pepperweed, Virginia	Star of Bethlehem	
Burclover	Fleabane, rough	(Peppergrass)	Strawberry, India mock	
Burdock	Galinsoga, hairy	Pigweed, prostate	Tansy ragwort	
Burweed, lawn	Galinsoga, smallflower	Pigweed, redroot	Tanweed	
(spurge)	Goldenrod	Pigweed, Russian	Thistle, blessed	
Buttercup, bulbous	Ground ivy	Pigweed, smooth	Thistle, bull	
Buttercup, creeping	Groundsel	Pigweed, tumble	Thistle, Canada	
Buttonweed, common	Gumweed	Pineapple weed	Thistle, Flodman's (prairie)	
(poorjoe)	Hairy fleabane	Plains coreopsis (tickseed)	Thistle, musk	
Buttonweed, Virginia	Hawkweed, meadow	Plantain, blackseed	Thistle, platte	
Carolina geranium (wild)	Hawkweed, mouseear	Plantain, bracted	Thistle, plumless	
Carpetweed	Hawkweed, orange	Plantain, broadleaf	Thistle, Russian	
Catnip	Hawkweed, yellow	Plantain, buckhorn	Thistle, scotch	
Catsear, spotted	Heal-all	Plantain, hoary	Thistle, wavyleaf	
(false dandelion)	Heartleaf drymary	Plantain, narrowleaf	Thistle, yellow (horrible)	
Chamber bitter	Hemp	Plantain, slender	Trailing crownvetch	
Chickweed, common	Henbit	Plantain, woolly	Velvetleaf	
Chickweed, mouseear	Hoary cress	Poison ivy	Vervain, blue	
Chickweed, sticky	Horsenettle	Poison oak	Vervain, hoary	
Chicory	Horseweed	Pokeweed	Vervain, prostrate	
Cinquefoil	Jimsonweed	Prairie Sunflower	Vervain, tall	
Clover, crimson	Johnny-Jumpup violet	Prickly lettuce	Vetch, bird	
Clover, Alyce	Knawel	(compass plant)	Vetch, hairy	
Clover, hop	Knotweed, prostrate	Prickly sida	Vetch, milk	
Clover, rabbit foot	Kochia	Prostrate knotweed	Vetch, narrowleaf	
Clover, red	Lambsquarters	Puncturevine	Vetch, wild	
Clover, strawberry	Lespedeza	Purslane, common	Virginia creeper	
Clover, yellow sweet	Mallow, alkali	Pusley, Brazil	Western clematis	
Clover, white sweet	Mallow, bristly	Pusley, Florida	Western salsify	
Clover, white	Mallow, common	Pusley, large flower	Wild aster	
Cockle	Mallow, dwarf	Ragweed, bur	Wild buckwheat	
Cocklebur	Mallow, Venice	Ragweed, common	Wild carrot	
Cocklebur, oriental	Marcela	Ragweed, lancleaf	Wild four-o'clock	

Spreader Settings

Spreader Directions:
Determining Your Lawn's Square Footage



To determine how much of this product to apply, you must determine the square footage of your lawn. An easy way to do this is to walk the length of your lawn, figuring that one pace equals about 3 feet. Do the same with the width of the lawn. Then multiply the length by the width to arrive at the total. Make sure you subtract the square footage of your home and driveway when calculating the total square footage of your lawn. If your lawn has an irregular shape, try dividing it up into a number of square areas. Calculate the square footage for each area, then add the figures together to reach your total lawn size.

Spreader Tips


The two main types of spreaders are the Drop Spreader, which drops the material straight down the width of the spreader as you push, and the Broadcast Spreader (also known as a rotary), which throws the material from the spreader.

Broadcast Spreader	Setting 2.6 Lbs./1,000 Sq. Ft.	Drop Spreader	Setting 2.6 Lbs./1,000 Sq. Ft.
Agri-Fab	4	Earthway	14
Earthway EV-N Spred	15	Republic	4
Republic EZGro	6½	Scotts AccuGreen	3½
Scotts Deluxe Edgeguard	4	Sta-Green	3½
Scotts EasyGreen	23		
Sta-Green	3		




These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Broadcast Spreader: Apply in a single pass by starting on the outside border and apply in a circular path working toward the center.



Drop Spreader: First make two passes along the top of the area you'll be covering. Then make two passes along the bottom area. Then move back and forth along the long part of the lawn. On each pass, as you reach the strips at the top and bottom of the area to be covered, shut off the spreader. It's also important to shut off the spreader when you are stopped.



Note: if spreader model does not appear above please call 800-233-1067, 8 am - 5 pm ET, Monday - Friday for additional model settings.

The above spreader settings are approximate. To check your spreader: 1 lb. of material should cover approximately 385 sq. ft. (2.6 lbs. of material per 1,000 sq. ft.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.
Pesticide Storage: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.
Pesticide Disposal and Container Handling: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Questions or Comments? Call 1-800-233-1067.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

**27-0-4
GUARANTEED ANALYSIS**

Total Nitrogen (N)	27%
3.4% Ammoniacal Nitrogen	
2.2% Water Insoluble Nitrogen*	
18.0% Urea Nitrogen	
3.4% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	4%
Sulfur (S)	3.8%
3.8% Combined Sulfur (S)	
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	4.0%
* 5.6% Slowly Available Nitrogen from Methylene Ureas.	

F699

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>.



Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your problem and simply send proof of purchase to Lebanon Seaboard Corporation, 1600 East Cumberland Street, Lebanon, PA 17042.

•For more information on our complete line of GreenView[®] Lawn and Garden Products, visit our website at www.greenviewfertilizer.com
•GreenView and GREENSMART are registered trademarks of Lebanon Seaboard Corporation.



21-31170

i/24

EPA Reg. No. 961-415

(L) (D)
EPA Est. No. 961-PA-01, 961-IL-01
Superscript used is first letter of lot number.