

GreenView®
Premium Formula

Spring Fertilizer
WEED & FEED
AND CRABGRASS PREVENTER

Covers 10,000 Sq. Ft.

24-0-6

NET WT.: 36 LBS. (16.3 KG)



GreenView® Premium Formula

Spring Fertilizer WEED & FEED AND CRABGRASS PREVENTER

Kills Lawn Weeds - Crabgrass,
Dandelions, Viney Weeds, Even Clover

Extended Time-Release Feeding
Of Nitrogen Lasts Into The Summer
Guaranteed Satisfaction

KEEP OUT OF REACH
OF CHILDREN
CAUTION

NOTICE: Read the entire label. Use only
according to label directions.

See back panel for complete Precautionary
Statements, including First Aid and
Directions for Use, including Storage and
Disposal.


ACTIVE INGREDIENTS:

2,4-D 0.04%
Mecoprop-p 0.14%
Dicamba 0.09%
Glycophosate 99.00%
OTHER INGREDIENTS: 100.00%

THIS PRODUCT CONTAINS:
12.5 lb 2,4-dichlorophenoxyacetic acid
equivalent per ton.
2.8 lb (2-(2-methyl-4-chlorophenoxy)
propionic acid equivalent per ton
1.2 lb 3,6-dichloro-*o*-benzoic acid equivalent
per ton.
52 lb diethothyper for
Inher Specific by AOC Methods
EPA Reg. No. 2217-989-961
EPA Est. No. 961-PA-01, 961-IL-01
Superscript used is first letter of lot number.

0 88685 30045 5

Spring Fertilizer WEED & FEED AND CRABGRASS PREVENTER



61-00045
0 88685 30045 5

GreenView®
Premium Formula

Spring Fertilizer WEED & FEED AND CRABGRASS PREVENTER

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION
Causes serious eye damage if it comes into contact with eyes. If it comes into contact with eyes, flush immediately with water for 15-20 minutes. Avoid contact with skin, eyes or clothing. Wash long-sleeved shirt and pants, gloves, socks, and waterproof gloves. Wash thoroughly after handling and before eating, drinking, chewing gum, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID
IF IN EYES:
• Rinse eye open and rinse slowly and gently with water for 15 - 20 minutes. Continue rinsing eye.
• Call a poison control center or doctor immediately for treatment advice.

IF SWALLOWED:
• If person is conscious, give a glass of water to swallow.
• Do not induce vomiting unless told to do so by a poison control center or doctor.

IF ON SKIN:
• Wash skin immediately with plenty of water for 15 - 20 minutes.
• Call a poison control center or doctor for treatment advice.

IF INHALED:
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
• Call a poison control center or doctor for treatment advice.

How to treat a person for a poison control center or doctor when you call: Application around a cistem or well may result in contamination of drinking water or groundwater.

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or off into storm drains, drainage ditches, garden surface waters. Applying this product in calm weather when it is recommended for the next 24 hours will help prevent wind or rain from washing pesticides off the treated area. Sweeping any products that land on driveways, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run-off into water bodies or drainage systems.

This chemical's properties and characteristics associated with chemicals detected in groundwater. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistem or well may result in contamination of drinking water or groundwater.

PREVENTIONAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or off into storm drains, drainage ditches, garden surface waters. Applying this product in calm weather when it is recommended for the next 24 hours will help prevent wind or rain from washing pesticides off the treated area. Sweeping any products that land on driveways, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run-off into water bodies or drainage systems.

Chemical's properties and characteristics associated with chemicals detected in groundwater.

Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistem or well may result in contamination of drinking water or groundwater.

WEEDS CONTROLLED

When applied as directed, this product will control the following weeds:

Annual Bluegrass: Creeping Bentgrass: Kikuyugrass (seeding)

Annual ryegrass: Croftgrass: Perennial Ryegrass: Smutgrass (seeding)

Barnyardgrass: Foxtail (green and yellow): Gooseneck

Established Cool Season Turfgrasses - Kentucky bluegrass, perennial ryegrass, fine and tall fescue

Established Warm Season Turfgrasses - Bahiagrass, bermudagrass, centipedegrass, Johnson grass, St. Augustine grass except "Floratam". Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established roots.

Do not use this product on seedling buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established roots.

Established Warm Season Turfgrasses - Colonial bentgrass, dichondra, carpetgrass, or "Floratam" St. Augustinegrass.

If the lawn has been seed treated, wait until after the third mowing before using this product. Do not seed, overseed or spray the treated area for 10 weeks after application.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

HOW MUCH TO USE:

Apply 3.6 pounds of product per 1,000 sq. ft. of lawn.

In the Fall: Scatter Do not exceed one application of the recommended rate of 3.6 lb of product per 1,000 sq. ft. (0.25 lb diethothyper per acre).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq. ft. per application (1.0 lb 2,4-D ae, 0.22 lb MCP-p ae, 0.09 lb dicamba (2.0 lb 2,4-D ae, 0.44 lb MCP-p ae and 0.18 lb dicamba as and 0.50 lb diethothyper per acre).

Applying at a rate higher than the maximum broadcast application rate may result in damage to plants.

Applying at a rate lower than the maximum broadcast application rate may result in poor results.

Overlap the coverage slightly, and shut off the spreader before making hard turns.

BROADCAST SPREADERS:

Scotts Turf Builder EdgeGuard..... 3.75

Scotts Turf Builder Hand-Powered..... 4

Sta-Green 220 Precision..... 4.25

Agri-Fab 45-0463 130 ft. Tow-Behind..... 3.5

DROP SPREADERS:

Scotts Turf Builder Classic Drop..... 5

Yardworks 750..... 3.25

Agri-Fab 175 ft Tow Behind 45-0288..... 3.5

If your spreader model is not listed, go to our website at www.greenviewfertilizer.com or call toll-free 800-233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model.

Storage and Disposal:

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food, feed, seeds or other pesticides.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly used: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER: FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, WITH RESPECT TO THIS PRODUCT, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY STATEMENT cannot be varied by any written or verbal statements or agreements at the point of sale or distribution.

Lebanon Seaboard Corporation guarantees you will be satisfied with the performance of this product when used as directed, or your money back. If not satisfied, describe the nature of your return and simple send proof of purchase to Lebanon Seaboard Corporation, 1600 E. Cumberland Street, Lebanon, PA 17042.

GreenView® is a registered trademark of Lebanon Seaboard Corporation.


MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 E. CUMBERLAND STREET
LEBANON, PA 17042

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lebsea.com>.

WHEN TO APPLY:
GREENVIEW PREMIUM FORMULA SPRING FERTILIZER WEED & FEED and CRABGRASS PREVENTER can be used either in the spring or fall. GREENVIEW PREMIUM FORMULA SPRING FERTILIZER WEED & FEED and CRABGRASS PREVENTER will control your lawn's weeds in one application. The grass and weeds are actively growing and the soil is moist. Do not apply when the grass and weeds are not growing.

IN THE FALL: The ideal time to apply GREENVIEW PREMIUM FORMULA SPRING FERTILIZER WEED & FEED and CRABGRASS PREVENTER is after the forsythia has bloomed or when the first dandelion flowers appear. In southern areas (see map), make a second application of GREENVIEW PREMIUM FORMULA SPRING FERTILIZER WEED & FEED and CRABGRASS PREVENTER 6 to 8 weeks after the first application.

IN THE SPRING: apply GREENVIEW PREMIUM FORMULA SPRING FERTILIZER WEED & FEED and CRABGRASS PREVENTER during the weather cool and both the grass and weeds are actively growing, generally around Labor Day.



HOW TO APPLY:
This product needs to stick to the leaves of broadleaf weeds and should be applied directly to the leaves. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor coverage or water runoff may result.

HOW MUCH TO USE:
Apply 3.6 pounds of product per 1,000 sq. ft. of lawn.

In the Fall: Scatter Do not exceed one application of the recommended rate of 3.6 lb of product per 1,000 sq. ft. (0.25 lb diethothyper per acre).

Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq. ft. per application (1.0 lb 2,4-D ae, 0.22 lb MCP-p ae, 0.09 lb dicamba (2.0 lb 2,4-D ae, 0.44 lb MCP-p ae and 0.18 lb dicamba as and 0.50 lb diethothyper per acre).

Applying at a rate higher than the maximum broadcast application rate may result in damage to plants.

Applying at a rate lower than the maximum broadcast application rate may result in poor results.

Overlap the coverage slightly, and shut off the spreader before making hard turns.

BROADCAST SPREADERS:

Scotts Turf Builder EdgeGuard..... 3.75

Scotts Turf Builder Hand-Powered..... 4

Sta-Green 220 Precision..... 4.25

Agri-Fab 45-0463 130 ft. Tow-Behind..... 3.5

DROP SPREADERS:

Scotts Turf Builder Classic Drop..... 5

Yardworks 750..... 3.25

Agri-Fab 175 ft Tow Behind 45-0288..... 3.5

If your spreader model is not listed, go to our website at www.greenviewfertilizer.com or call toll-free 800-233-1067, 8 A.M. - 5 P.M. EDT, Monday - Friday, for additional model.

www.greenviewfertilizer.com



61-00045
0 88685 30045 5

GreenView®
Premium Formula

Spring Fertilizer WEED & FEED AND CRABGRASS PREVENTER