

A.C.R.E. acidifying fertilizer

PROFESSIONAL LANDSCAPE NUTRITION

Lebanon Seaboard Corporation1600 East Cumberland Street • Lebanon, PA 17042
www.lebanonturf.com

NET WT. 50 LBS. (22.7 KG)



PROFESSIONAL LANDSCAPE NUTRITION

Total Nitrogen (N)	12%	Total Nitrogen (N)	12%
4.6% Ammoniacal Nitrogen		4.6% Ammoniacal Nitrogen	
4.8% Water Insoluble Nitrogen*		4.8% Water Insoluble Nitrogen*	
0.3% Urea Nitrogen		0.3% Urea Nitrogen	
2.3% Other Water Soluble Nitrogen*	221	2.3% Other Water Soluble Nitrogen*	00/
Available Phosphate (P ₂ O ₅)	3%	Available Phosphate (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	6%	Soluble Potash (K ₂ O)	6%
Sulfur (S)	4.6%	Sulfur (S)	4.6%
Iron (Fe)	2.5%	Iron (Fe) (actual)	2.5%
Manganese (Mn)	0.5%	Manganese (Mn) (actual)	0.5%
0.06% Water Soluble Manganese (Mn)	•	0.06% Water Soluble Manganese (Mn)	•
Derived from: Ammonium Phosphate, Ammonium S		Chlorine (Cl) (maximum)	5.0%
Biosolids, Methylene Ureas, Urea, Muriate of Potash Oxide, Ferrous Sulfate, Manganese Sucrate.	ո, Ferric	Derived from: Ammonium Phosphate, Ammonium S Biosolids, Methylene Ureas, Urea, Muriate of Potash	
*7.1% Slowly Available Nitrogen from Methylene Ureas.		Oxide, Ferrous Sulfate, Manganese Sucrate.	,
Chlorine (CI) not more than	5.0%	*7.1% Slowly Available Nitrogen from Methylene Ure	eas.
	U.S.A.	-	Cana

general

Woodace A.C.R.E. 12-3-6 is a premium fertilizer for maintaining foundation plantings, shrubs, ground covers, and trees that prefer a slightly acidic soil environment. Woodace A.C.R.E. 12-3-6 is also ideal for fertilizing plants where micronutrient availability is limited by high soil pH. Woodace A.C.R.E. will aid in maintaining a slightly acid soil pH which is ideal for most evergreens and the majority of foundation plants. Additional Iron is included to help enhance the color and appearance of foliage.

application

Woodace A.C.R.E. is designed for top-dress application to ornamentals growing in planting beds and other non-turf areas. **For individual plants** apply Woodace A.C.R.E. evenly within the dripline or known root area. The feeder roots of large shrubs and trees may extend well beyond the dripline. Adjust the area accordingly. **For densely planted areas or ground covers** broadcast Woodace A.C.R.E. evenly over the entire area at a rate of 0.8 - 2.5 lbs. per 100 sq. ft., 1 - 3 lbs. N/1000 sq. ft. (0.4 kg - 1.1kg per 9.3 m², 0.45 - 1.36 kg.N/92.9m²). Adjust the height and trajectory of the spreader where possible to minimize reflecting granules off of dense foliage. Do not broadcast fertilizer onto wet foliage. Fertilizer should be applied when plants are actively growing. Do not attempt to incorporate fertilizers into the root area of established plants. Never fertilize when plants are suffering from stress related to transplant, moisture, heat, insect, disease or abiotic injury. Generally spring and fall provide the optimum time to apply supplemental nutrition.

fertilization program guidelines

Do not wash off with water.

Woodace A.C.R.E. should be a part of an integrated landscape maintenance program. In addition to fertilization, pest monitoring, weed control, pruning, and water management are essential. In most landscape situations nutrients are not naturally replenished at a sufficient rate to guarantee optimum health and appearance. The objective of most landscape fertilization programs is to improve the health and appearance of plantings by providing proper fertility. Other objectives for which Woodace A.C.R.E. may be used include: supplementing nutrients known to be deficient, encouraging foliar or root growth, increasing bloom or fruiting, or to aid in recovery from injury or disease. These label directions are intended to serve as a general guideline for the use of Woodace A.C.R.E. Other resources such as soil testing, local extension recommendations, and specific plant knowledge should be incorporated whenever possible.

WARNING

HAZARD STATEMENTS

Harmful if swallowed.
Causes skin irritation.
May be harmful if inhaled.
May cause respiratory irritation.

Causes eye irritation.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands on impervious surfaces back into your turf.

PRECAUTIONARY STATEMENTS

Wash hands and exposed skin thoroughly after handling.

Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection.

If in eyes, rinse with water and seek medical attention if symptoms persist.

Rinse mouth.

Take off contaminated clothing and wash before reuse.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell.

If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Store locked up.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container properly as required by law for waste pesticides. Keep out of reach of children

Precautionary Statements for disposal - Dispose in accordance with all federal, state and loca regulations.

rate chart

broadcast (for heavily planted beds)

N rate Ib	s./100 sq. ft. (kg/9.3m²)	lbs./1000 sq. ft. (kg/92.9m²)
1 lb./1000 ft ² (0.45 kg/92.9m ²)	0.8 lb. (0.4 kg)	8.3 lbs. (3.8 kg)
2 lbs./1000 ft² (0.9 kg/92.9m²)	1.6 lbs. (0.7 kg)	17 lbs. (7.7 kg)
3 lbs./1000 ft² (1.4 kg/92.9m²)	2.5 lbs. (1.1 kg)	25 lbs. (11.3 kg)

shrubs (per plant within dripline)

(por plant than anymio)					
size	rate	measure			
8-12" (20-30 cm)	2 oz.	1⁄4 cup			
13-18" (30-46 cm)	4 oz.	½ cup			
19-24" (48-61 cm)	6 oz.	3∕4 cup			
25-30" (64-76 cm)	8 oz.	1 cup			
31-36" (79-91 cm)	12 oz.	1½ cups			
37-42" (94-107 cm)	16 oz.	2 cups			
> 42" (> 107 cm)	1 cup per foot (0.3 meters)				
	of branch spread				

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.



Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042

800-233-0628 • (717) 273-1685 Made in the U.S.A. www.LebanonTurf.com

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-lebsea.com

One cup is approximately 8 oz. (227 grams) by weight.

Shrub size refers to branch spread for shrubs where height and spread are approximately the same. For fastigate or decumbent forms reduce the amount by approximately ½.

The user assumes all liability for applications of this product. For uses that are not specifically listed on the product label, it is recommended that the user perform a test on a limited basis before commencing large scale applications.

Woodace® is a registered trademark of Lebanon Seaboard Corporation.





F699